

Palladium-Catalyzed Insertion of CO₂ into Vinylaziridines: New Route to 5-Vinyloxazolidinones


Francesco Fontana, C. Chun Chen, and Varinder K. Aggarwal*

School of Chemistry, University of Bristol, Cantock's Close, Bristol, U.K.

V.Aggarwal@bristol.ac.uk

Received May 5, 2011

ABSTRACT



2-Vinylaziridines undergo a mild Pd-catalyzed ring-opening cyclization reaction with an ambient atmosphere of carbon dioxide to give 5-vinyloxazolidinones. The process is high yielding as well as regio- and stereoselective.

Vinylaziridines are valuable synthetic intermediates¹ which, as an illustration, undergo a host of very useful palladium-catalyzed transformations (Scheme 1). For example, we² and others³ showed that they could be converted into unsaturated β -lactams with a high level of stereocontrol. In the presence of doubly activated Michael acceptors they can also be converted into pyrrolidines.^{4a,b} We recently extended this methodology to singly activated acceptors and demonstrated its utility in a stereocontrolled synthesis of (+)-kainic acid.^{4c} The Pd-catalyzed reaction of vinylaziridines with heterocumulenes

(e.g., isocyanates, carbodiimides, and isothiocyanates) is also well established,⁵ but insertion into the archetypal heterocumulene, CO₂, has not been reported.⁶ Such a reaction would give vinyloxazolidinones, important motifs in both synthesis^{7–9} and pharmacology.¹⁰ However, this reaction did not seem promising since Ibuka had reported that 3-tosyl-5-vinyloxazolidin-2-ones underwent Pd-catalyzed *decarboxylation* to give *cis*-*N*-tosyl-2-vinylaziridines (Scheme 1).¹¹ Furthermore, Knight^{4b,12} had shown that vinyloxazolidinones bearing electron-withdrawing groups

(1) For reviews, see: (a) Ohno, H. In *Aziridines and Epoxides in Asymmetric Synthesis*; Yudin, A. K., Ed.; Wiley-VCH: Weinheim, Germany, 2006; Chapter 2. (b) Sweeney, J. B. In *Science of Synthesis*; Enders, D., Ed.; Georg Thieme Verlag: 2009; Vol. 40a, pp 643–772. (c) Müller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905–2920. (d) Sweeney, J. B. *Aziridines and Epoxides in Asymmetric Synthesis*; Yudin, A. K., Ed.; Wiley-VCH: Weinheim, Germany, 2006; Chapter 4.


(2) Fontana, F.; Tron, G. C.; Barbero, N.; Ferrini, S.; Thomas, S. P.; Aggarwal, V. K. *Chem. Commun.* **2010**, *46*, 267–269.

(3) (a) Tanner, D.; Somfai, P. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 2415–2418. (b) Spears, G. W.; Nakanishi, K.; Ohfuné, Y. *Synlett* **1991**, 91–92.

(4) (a) Aoyagi, K.; Nakamura, H.; Yamamoto, Y. *J. Org. Chem.* **2002**, *67*, 5977–5980. (b) For an analogous reaction on vinyloxazolidinones, see: Knight, J. G.; Stoker, P. A.; Tchabanenko, K.; Harwood, S. J.; Lawrie, K. W. M. *Tetrahedron* **2008**, *64*, 3744–3750. (c) Lowe, M. A.; Ostovar, M.; Ferrini, S.; Chen, C. C.; Lawrence, P. G.; Fontana, F.; Calabrese, A. A.; Aggarwal, V. K. *Angew. Chem., Int. Ed.* **2011**, DOI: 10.1002/anie.201101389.

(5) For selected examples, see: (a) Butler, D. C. D.; Inman, G. A.; Alper, H. *J. Org. Chem.* **2000**, *65*, 5887–5890. (b) Fandrick, D. R.; Trost, B. M. *J. Am. Chem. Soc.* **2003**, *125*, 11836–11837. (c) Dong, C.; Alper, H. *Tetrahedron: Asymmetry* **2004**, *15*, 1537–1540.


Scheme 1. Pd-Catalyzed Applications of Vinylaziridines



on nitrogen readily lost CO₂ upon treatment with Pd(0) and that the resulting π -allyl Pd intermediate could be trapped with CO or dipolarophiles.

Thus, the potential for incorporating CO₂ into vinylaziridines,¹³ perhaps because of its high thermodynamic and kinetic stability, seemed remote. Nevertheless, our interest in the synthesis¹⁴ and utility² of vinylaziridines together with the contemporary environmental need to find further uses of CO₂ as a raw material^{15,16} prompted us to investigate this area further.

Scheme 2. Synthesis of Vinylaziridines



We initially synthesized a series of *N*-tosyl-2-vinylaziridines **1** using our sulfur ylide methodology as shown in Scheme 2 (see Supporting Information (SI)).² Treatment of the vinylaziridine **1a** with Pd₂(dba)₃·CHCl₃ in the presence of Ph₃P and just an *ambient* atmosphere of CO₂ gave the *trans*-vinylloxazolidinone **3a** in a promising 61% yield. In addition, *cis*-aziridine **2a** was also obtained together with traces of the starting material, *trans*-aziridine **1a** (Table 1, entry 1).¹⁷

(6) Trost has reported the insertion of CO₂ into vinyl epoxides at elevated pressure. See: (a) Trost, B. M.; Angle, S. R. *J. Am. Chem. Soc.* **1985**, *107*, 6123–6124. Yoshida reported the conversion of aminoalkenyl carbamates into vinylloxazolidinones. See: (b) Yoshida, M.; Ohsawa, Y.; Sugimoto, K.; Hidetoshi, T.; Ihara, M. *Tetrahedron Lett.* **2007**, *48*, 8678–8682.

(7) (a) For a recent review, see: Zappia, G.; Cancelliere, G.; Gacs-Baitz, E.; Delle Monache, G.; Misiti, D.; Nevola, L.; Botta, B. *Curr. Org. Synth.* **2007**, *4*, 238–307.

(8) For representative examples on the use of 5-vinylloxazolidinones in synthesis, see: (a) Reference 4b. (b) Cook, G. R.; Shanker, P. S.; Peterson, S. L. *Org. Lett.* **1999**, *1*, 615–617. (c) Yamamoto, T.; Hasegawa, H.; Ishii, S.; Kaji, S.; Masuyama, T.; Harada, S.; Katsumura, S. *Tetrahedron* **2008**, *64*, 11647–11660. (d) Robertson, J.; Abdulmalek, E. *Tetrahedron Lett.* **2009**, *50*, 3516–3518.

(9) For examples of hydrolysis of vinylloxazolidinones to unsaturated β -aminoalcohols, see: (a) Olofsson, B.; Khamrai, U.; Somfai, P. *Org. Lett.* **2000**, *2*, 4087–4089. (b) Disadee, W.; Ishikawa, T. *J. Org. Chem.* **2005**, *70*, 9399–9406.

(10) (a) Colca, J. R.; McDonald, W. G.; Waldon, D. J.; Thomasco, L. M.; Gadwood, R. C.; Lund, E. T.; Cavey, G. S.; Mathews, W. R.; Adams, L. D.; Cecil, E. T.; Pearson, J. D.; Bock, J. H.; Mott, J. E.; Shinabarger, D. L.; Xiong, L.; Mankin, A. S. *J. Biol. Chem.* **2003**, *278*, 21972–21979. (b) Hoellman, D. B.; Lin, G.; Ednie, L. M.; Rattan, A.; Jacobs, M. R.; Appelbaum, P. C. *Antimicrob. Agents Chemother.* **2003**, *47*, 1148–1150. (c) Jacqueline, C.; Caillon, J.; Le Mabeccque, V.; Miegerville, A.-F.; Donnio, P.-Y.; Bugnon, D.; Potel, G. *J. Antimicrob. Chemother.* **2003**, *51*, 857–864. (d) Moellering, R. C., Jr. *Ann. Intern. Med.* **2003**, *138*, 135–142. (e) Rubinstein, E.; Isturiz, R.; Standiford, H. C.; Smith, L. G.; Oliphant, T. H.; Cammarata, S.; Hafkin, B.; Le, V.; Remington, J. *Antimicrob. Agents Chemother.* **2003**, *47*, 1824–1831. (f) Mansour-Ghanaei, F.; Fallah, M. S.; Shafaghi, A. *Med. Sci. Monit.* **2002**, *8*, 127–130. (g) Kaatz, G. W.; Rybak, M. J. *Emerging Drugs* **2001**, *6*, 43–55. (h) Fluit, A. C.; Schmitz, F. J.; Verhoef, J.; Milatovic, D. *J. Antimicrob. Chemother.* **2002**, *50*, 271–276.

The formation of the *cis*-aziridine **2a** is in keeping with Ibuka's observations as he had previously found that *trans*-*N*-tosyl aziridines readily isomerized to *cis*-aziridines (98:2) in the presence of Pd(0).¹¹

The isolation of the isomerized starting material **2a** implicated a slow reaction between the amide anion and CO₂. We reasoned that ion pairing of the amide anion with the cationic Pd complex might compromise its reactivity and that increased nucleophilicity of the aza-anion could be achieved by rendering the Pd center neutral. Such a change had been pivotal in the successful reactions of vinyl aziridines with singly activated Michael acceptors.^{4c} We therefore decided to add tetrabutylammonium chloride (TBAC).¹⁸ While reaction at RT was lower yielding, at 0 °C the product vinylloxazolidinone **3a** was obtained in 86% yield (entries 2 and 3). We were able to achieve similar results at a lower phosphine loading (entry 4). Replacing the highly hygroscopic TBAC with the more easily manageable fluoride source tetrabutylammonium difluorotriphenylsilicate (TBAT) led to a further increase in yield (92%) with no detectable traces of starting material **1a** or *cis*-aziridine **2a**. Furthermore the reaction was now completed over a shorter reaction time (entry 5). A control experiment in the absence of TBAT or Pd (entries 6, 7) confirmed the beneficial effects of both species in promoting the carboxylation.

Having developed an optimized protocol we then screened the scope of the reaction (Table 2). Carboxylation occurred uneventfully for a series of diaryl-substituted *trans*-aziridines **1b–1g** bearing either electron-donating or -withdrawing *para* substituents (entries 2, 3, 6, 7, 9). However, when R¹ possessed greater electron-donating character the addition of TBAT became detrimental and

(11) Ibuka, T.; Mimura, N.; Aoyama, H.; Akaji, M.; Ohno, H.; Miwa, Y.; Taga, T.; Nakai, K.; Tamamura, H.; Fujii, N. *J. Org. Chem.* **1997**, *62*, 999–1015.

(12) (a) Knight, J. G.; Ainge, S. W.; Harm, A. M.; Harwood, S. J.; Maughan, H. I.; Armour, D. R.; Hollinshead, D. M.; Jaxa-Chamiec, A. A. *J. Am. Chem. Soc.* **2000**, *122*, 2944–2945. (b) Knight, J. G.; Tchabanenko, K. *Tetrahedron* **2002**, *58*, 6659–6664. (c) Knight, J. G.; Lawson, I. M.; Johnson, C. N. *Synthesis* **2006**, 227–230.

(13) For examples of alkyl aziridines into oxazolidinones catalyzed by LiI, see: (a) Miller, A. W.; Nguyen, S. T. *Org. Lett.* **2004**, *6*, 2301–2304. (b) Hancock, M. T.; Pinhas, A. R. *Synthesis* **2004**, *14*, 2347–2355. (c) Mu, W.-H.; Chasse, G. A.; Fang, D.-C. *J. Phys. Chem. A* **2008**, *112*, 6708–6714.


(14) (a) Aggarwal, V. K.; Alonso, E.; Fang, G.; Ferrara, M.; Hynd, G.; Porcelloni, M. *Angew. Chem., Int. Ed.* **2001**, *40*, 1433–1436. (b) Illa, O.; Arshad, M.; Ros, A.; McGarrigle, E. M.; Aggarwal, V. K. *J. Am. Chem. Soc.* **2010**, *132*, 1828–1830. For a recent review, see: (c) Li, A.-H.; Dai, L.-X.; Hou, X.-L.; Chen, M.-B. *J. Org. Chem.* **1996**, *61*, 4641–4648. (d) Zhu, B.-H.; Zheng, J.-C.; Yu, C.-B.; Sun, X.-L.; Zhou, Y.-G.; Shen, Q.; Tang, Y. *Org. Lett.* **2010**, *3*, 504–507. (e) McGarrigle, E. M.; Myers, E. L.; Illa, O.; Shaw, M. A.; Riches, S. L.; Aggarwal, V. K. *Chem. Rev.* **2007**, *107*, 5841–5883. For routes from aziridine aldehydes, see: (f) Baktharaman, S.; Afagh, N.; Vandersteen, A.; Yudin, A. K. *Org. Lett.* **2010**, *12*, 240–243.

(15) (10) For a recent review on the use of CO₂ in synthesis, see: Sakakura, T.; Choi, J.-C.; Yasuda, H. *Chem. Rev.* **2007**, *107*, 2365–2387.

(16) For an example of waste CO₂ being converted into useful molecules, see: North, M.; Villuendas, P.; Young, C. *Chem.—Eur. J.* **2009**, *15*, 11454–11457.


(17) To avoid any further reaction of the desired oxazolidinone, the reaction mixture was promptly filtered through a pad of silica to remove the catalyst (see Supporting Information).

(18) For a review on the effects of halide additives in palladium chemistry, see: Fagnou, K.; Lautens, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 26–47.

Table 1. Optimization of the Carboxylation Reaction^a

entry	[Pd] mol %	PPh ₃ /Pd	additive (equiv)	<i>t</i> (°C)	time (h)	conversion (%) ^b	yield 2a (%) ^b	yield 3a (%) ^b
1	10	6	–	rt	8	98	29	61
2	10	6	TBAC (0.2)	rt	18	99	11	42
3	10	6	TBAC (0.2)	0	18	100	traces	86
4	10	2	TBAC (0.2)	0	4	100	–	86
5	10	2	TBAT (0.2)	0	2	100	–	92^c
6	10	2	–	0	2	98	–	71
7 ^d	–	–	TBAT (0.2)	0	2	0 ^e	–	–

^a dba = dibenzylideneacetone. TBAC = tetrabutylammonium chloride. TBAT = tetrabutylammonium difluorotriphenylsilicate. Aziridine concentration = 6×10^{-2} M. ^b Determined by ¹H NMR using 1,1,2,2-tetrachloroethane as an internal standard. ^c Isolated yield. ^d Carried out in the presence of PPh₃ (0.2 equiv). ^e Only starting material was recovered.

Table 2. Scope of Pd-Catalyzed Carboxylation Reaction^a

entry	aziridine	R ¹	R ²	<i>dr</i> (1:2)	time (h)	yield (%) ^b	<i>dr</i> (3:4) ^c
1	1a	Ph	Ph	>98:2	2	92	>98:2
2	1b	Ph	<i>p</i> -Me-C ₆ H ₄	>98:2	2	86	>98:2
3	1c	Ph	<i>p</i> -OMe-C ₆ H ₄	97:3	4	87	97:3
4	1c/2c	Ph	<i>p</i> -OMe-C ₆ H ₄	52:48	4	62 ^d	79:21
5	2c	Ph	<i>p</i> -OMe-C ₆ H ₄	2:>98	4	40	8:92
6	1d	Ph	<i>p</i> -Cl-C ₆ H ₄	>98:2	4	86	>98:2
7	1e	<i>p</i> -Me-C ₆ H ₄	Ph	>98:2	4	89	>98:2
8 ^e	1f	<i>p</i> -OMe-C ₆ H ₄	Ph	>98:2	4	91	>98:2
9	1g	<i>p</i> -Cl-C ₆ H ₄	Ph	>98:2	2	84	>98:2
10	1h	Ph	Me	>98:2	3	78	96:4
11	1i	Ph	TMS	>98:2	24	40	>98:2
12 ^e	1j	Cy	Ph	>98:2	4	83	>98:2
13	1k/2k	Ph	H	85:15	0.5	71 ^f	85:15
14 ^e	1l	H	H	–	0.5	68	–
15	(<i>2R,3R</i>)- 1a (98:2 <i>er</i>)	Ph	Ph	>98:2	2	89 (>99:1 <i>er</i>) ^g	>98:2


^a dba = dibenzylideneacetone. TBAT = tetrabutylammonium difluorotriphenylsilicate. Aziridine concentration = 6×10^{-2} M. ^b Isolated yield. ^c Determined by ¹H NMR of the crude. ^d Determined by ¹H NMR with 1,1,2,2-tetrachloroethane as internal standard. ^e Conducted in the absence of TBAT. ^f ¹H NMR of the crude showed the desired oxazolidinones together with *cis*-aziridine **2j** in a ratio of 85:15. ^g Determined by chiral HPLC (see S1).

higher yields were achieved without it (entry 8). Alkyl residues were well tolerated on both the vinyl moiety, **1h** (R² = Me), **1i** (R² = SiMe₃), and the aziridine core, **1i** (R¹ = Cy) (entries 10, 11, 12). The unsubstituted aziridine **1l** underwent carboxylation, but as before, higher yields were observed in the absence of TBAT (entry 13). Interestingly, the ratio of *trans*- and *cis*-vinylloxazolidinones reflected the ratio of *trans*- and *cis*-aziridines that we started with. Using enantioenriched aziridine (*2R,3R*)-**1a**, the *trans*-vinylloxazolidinone (+)-**3a** was obtained with

very high *er*, indicating that no erosion in enantiopurity took place (entry 14). We were also able to test the *cis*-aziridine **2c** and a ~1:1 mixture of *cis/trans* isomers **1c/2c** (entries 4, 5). Unexpectedly, the *cis*-aziridine mainly gave the *cis*-oxazolidinone **4c** and the ~1:1 mixture led to a 79:21 mixture of *trans/cis* oxazolidinones.

On the basis of these observations together with previous studies,² we propose the mechanism shown in Scheme 3. The *trans-E*-vinylaziridine can react with Pd(0) to give the π -allyl palladium intermediate **5**. Pd–Pd

Scheme 3. Proposed Mechanism




mediated isomerization¹⁹ would give the isomeric complex **6** which can ring close to the *cis-E*-vinylaziridine **2**. This is the accepted mechanism for isomerization of *trans*- and *cis*-vinylaziridines. Capture of the π -allyl palladium intermediates **5** and **6** by CO₂ then give the vinylloxazolidinones **3** and **4** respectively. The fact that *cis*-aziridine **2c** gives mostly the *cis*-oxazolidinone **4c** implies that isomerization of the π -allyl palladium intermediates **5** and **6** has been significantly suppressed under the reaction conditions. Factors that reduce the extent of such isomerization include (i) a lower temperature and (ii) use of a halide additive.^{20,4c} Indeed, at RT and in the absence of halide salts the *cis*-aziridine **2c** gave considerably more of *trans*-oxazolidinone (68 *cis*/32 *trans*) indicating that a greater degree of isomerization had taken place. Moreover, subjecting the *cis*-oxazolidinone **4c** (91 *cis*/9 *trans*) to the reaction conditions resulted in the same ratio of isomers indicating that no isomerization occurred. Again, at RT but in the absence of halide salts the isomerization of **4c** was much more rapid giving increased amounts of *trans*-oxazolidinone (76 *cis*/24 *trans*). These experiments show that *cis*–*trans* isomerization of both the aziridines and oxazolidinones is largely arrested under the reaction conditions (kinetic control) but can occur in the absence of halide ions and higher temperatures (thermodynamic control).

Knight has shown that 5-vinylloxazolidinones can serve as substrates for Pd(0)-catalyzed reactions, e.g. carbonylation, in a manner analogous to vinylaziridines.¹² However, subtle differences in product distributions are observed (Scheme 4). For example, *N*-tosyl-2-vinylaziridines where R² = H or Ar are known to afford β -lactams **9**.^{2,3a} However, Knight reported that submission of the corresponding *N*-tosyl-vinylloxazolidinones (R² = H) to carbonylation gave the δ -lactam derivative **10** instead.^{12c} The same regioisomers were obtained from vinylaziridines when

R² = SiMe₃.^{2,21} Surprisingly, we observed that carbonylation of vinylloxazolidinones **3a** (R¹, R² = Ph) and **3d** (R¹ = Ph, R² = *p*-Cl-C₆H₄) afforded the β -lactams **9**, albeit in lower yields and diastereoselectivity as compared to the corresponding aziridines (see SI for details).² This indicates that the intermediates involved prior to insertion of CO are subtly different. It is likely that decarboxylation is not instantaneous and that the product distribution is influenced by the rates of isomerization of the π -allyl Pd species and by the rates of carbonylation and decarboxylation, all of which will be substrate-dependent.

Scheme 4. Regioselectivity of Carbonylation



In conclusion, we have discovered that readily available *trans*-vinylaziridines can be converted into *trans*-vinylloxazolidinones through a Pd-catalyzed carbonylation process. The use of CO₂ in such processes is rare because CO₂ is widely regarded as thermodynamically stable and kinetically unreactive. The results presented herein show that it is far from kinetically unreactive, as should be expected for an electron-deficient carbon atom bearing two strongly activated electron sinks.

Acknowledgment. V.K.A. thanks the EPSRC for a Senior Research Fellowship. F.F. thanks “Università degli Studi di Milano” for funding. The authors thank Dr. Mairi Haddow (University of Bristol) for X-ray structure determination of compound **1i**.

Supporting Information Available. Synthesis and characterization of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(19) Gangberg, K. L.; Bäckvall, J.-E. *J. Am. Chem. Soc.* **1992**, *114*, 6858–6863.

(20) Cantat, T.; Agenet, N.; Jutand, A.; Pleixats, R.; Moreno-Manas, M. *Eur. J. Org. Chem.* **2005**, 4277–4286.

(21) We ascribed this result to an inherent preference for carbonylation to occur adjacent to Si due to the shorter C–Pd bond length. See: Branchadell, V.; Moreno-Manas, M.; Pleixats, R. *Organometallics* **2002**, *21*, 2407–2412.